

KIWIGREEN ACTION SHEET



If your Zespri KiwiGreen Monitoring round has found one or more of the below market access pests, the actions mentioned below must be taken to retain market access to China. Once action has been taken you must notify your PMC

**PESTS OF CONCERN FOR CHINA
REQUIRES ACTION FOR MARKET ACCESS**

| <u>PEST</u> | <u>FOUND</u> | <u>ACTION</u> | <u>PRODUCTS</u> | <u>REFERENCES</u> |
|---|--------------|---|---|--|
| <u>SCALE</u> | 4-7% | 1) Spray blocks above threshold and record spray in Spray Diary OR 2) Instruct PMC to remonitor blocks above threshold (prior to harvest) SPRAYS MUST BE RECORDED IN SPRAY DIARY WITH THE REASON <i>SCALE CONTROL</i> | EXCEL OB, EXCEL PLUS, DC TRON PLUS OB, ENSPRAY 99, | Using summer oils safely Monitoring and responding to scale in summer Armoured Scale Factsheet Summer Oil for Scale Control |
| | >7%-20% | 1) Spray blocks above threshold and record spray in Spray Diary AND THEN 2) Instruct PMC to re-monitor blocks above threshold (prior to harvest) SPRAYS MUST BE RECORDED IN SPRAY DIARY WITH THE REASON <i>SCALE CONTROL</i> | | |
| | >20% | 1) Spray blocks above threshold (2 sprays can be applied if timing is appropriate) and record spray in Spray Diary AND THEN 2) Instruct PMC to re-monitor blocks above threshold (prior to harvest) CN Hold will remain | | |
| EXPECTED ACTION IS VARIETY SPECIFIC. OIL ON G3 SAFE MID FEB ONLY. OIL ON GREEN SAFE JAN-END OF MARCH. OIL ON RED HAS NOT HAD SAFETY TESTED THEREFORE SPRAYING IS NOT EXPECTED | | | | |
| <u>LEAFROLLER</u> | 0.5% - 1% | 1) Spray blocks above threshold and record spray in Spray Diary SPRAYS MUST BE RECORDED IN SPRAY DIARY WITH THE REASON AS <i>LEAFROLLER CONTROL</i> | BIO-BIT DF, DELFIN WG, DIPEL DF, BACTUR, (PROCLAIM ETC IF 42 DAYS PRIOR TO HARVEST) | Leafroller Factsheet |
| | 1%-3% | 1) Spray blocks above threshold AND THEN 2) Instruct PMC to re-monitor blocks above threshold (prior to harvest) | | |
| | >3% | 1) Spray blocks and re-monitor. CN Hold will remain | | |

KIWIGREEN ACTION SHEET



| | | | | |
|-------------------------|--|---|--------------------------------------|--|
| <p><u>WHEAT BUG</u></p> | <p>LOAD OUT AREA NOT CLEAR OF HOST WEEDS</p> | <ol style="list-style-type: none"> 1) Clear high-risk loadout area of host weeds <ul style="list-style-type: none"> • A list of host weeds & images is shown in the attached Wheat Bug Fact Sheet. • A load out area is defined as any area, including a 3m perimeter, where bins will be placed awaiting transport to the packhouse. AND THEN 2) Record this in spray diary: <ul style="list-style-type: none"> • Select 'Add Spray line' and Enter details for a spray line <ol style="list-style-type: none"> i. In the notes section comment on the actions taken i.e. herbicide spraying / manual removal of host weeds for wheat bug ii. From the product drop down menu select either: Wheat bug management (Glyphosate) OR/ Wheat bug management (manual removal). iii. From the 'reason' dropdown select 'Wheat bug' iv. Under 'blocks' select 'loadout'. v. The remaining sections should be completed as placeholders if not relevant i.e. Spray contractor is not relevant if method of action is weed removal. <p>LOAD OUT AREA MUST REMAIN CLEARED UNTIL HARVEST</p> | <p>ZESPRI APPROVED HERBICIDE</p> | <p>Wheat Bug Factsheet</p> |
|-------------------------|--|---|--------------------------------------|--|